


# 2-Substituted pyrimidines

## Abstract

### 5 2-Substituted pyrimidines of the formula I



in which the index n and the substituents L, R<sup>a</sup>, R<sup>b</sup>, R<sup>c</sup>, R<sup>z</sup>, R<sup>u</sup>, R<sup>v</sup>, A', A'' and A''' are as defined in the description and:

10

R<sup>1</sup> is C<sub>3</sub>-C<sub>10</sub>-alkyl, C<sub>3</sub>-C<sub>10</sub>-alkenyl, C<sub>3</sub>-C<sub>10</sub>-alkynyl, C<sub>3</sub>-C<sub>12</sub>-cycloalkyl, C<sub>3</sub>-C<sub>10</sub>-cycloalkenyl or a five- to ten-membered saturated, partially unsaturated or aromatic heterocycle which is attached via carbon and contains one to four heteroatoms from the group consisting of O, N and S,

15

R<sup>2</sup> is halogen, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, C<sub>2</sub>-C<sub>4</sub>-alkynyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>3</sub>-C<sub>4</sub>-alkenyloxy or C<sub>3</sub>-C<sub>4</sub>-alkynyloxy, where the alkyl, alkenyl and alkynyl radicals of R<sup>2</sup> may be substituted by halogen, cyano, nitro, C<sub>1</sub>-C<sub>2</sub>-alkoxy or C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl,

20 and

R<sup>3</sup> is cyano, CO<sub>2</sub>R<sup>a</sup>, C(=O)NR<sup>z</sup>R<sup>b</sup>, C(=O)-N-OR<sup>b</sup>, C(=S)-NR<sup>a</sup>R<sup>b</sup>, C(=NOR<sup>a</sup>)NR<sup>z</sup>R<sup>b</sup>, C(=NR<sup>a</sup>)NR<sup>z</sup>R<sup>b</sup>, C(=O)NR<sup>a</sup>-NR<sup>z</sup>R<sup>b</sup>, C(=N-NR<sup>z</sup>R<sup>c</sup>)NR<sup>a</sup>R<sup>b</sup>, C(=O)R<sup>a</sup>, C(=NOR<sup>b</sup>)R<sup>a</sup>, C(=N-NR<sup>z</sup>R<sup>b</sup>)R<sup>a</sup>, CR<sup>a</sup>R<sup>b</sup>-OR<sup>z</sup>, CR<sup>a</sup>R<sup>b</sup>-NR<sup>z</sup>R<sup>c</sup>,  
 25 ON(=CR<sup>a</sup>R<sup>b</sup>), O-C(=O)R<sup>a</sup>,  
 NR<sup>a</sup>R<sup>b</sup>, NR<sup>a</sup>(C(=O)R<sup>b</sup>), NR<sup>a</sup>(C(=O)OR<sup>b</sup>), NR<sup>a</sup>(C(=O)-NR<sup>z</sup>R<sup>b</sup>), NR<sup>a</sup>(C(=NR<sup>c</sup>)R<sup>b</sup>),  
 NR<sup>a</sup>(N=CR<sup>c</sup>R<sup>b</sup>), NR<sup>a</sup>-NR<sup>z</sup>R<sup>b</sup>, NR<sup>z</sup>-OR<sup>a</sup>, NR<sup>a</sup>(C(=NR<sup>c</sup>)-NR<sup>z</sup>R<sup>b</sup>), NR<sup>a</sup>(C(=NOR<sup>c</sup>)R<sup>b</sup>),

30

processes for preparing these compounds, compositions comprising these compounds and their pesticidal use are described.